

## Article

# Monitoring for airborne respiratory viruses in a general pediatric ward in Singapore

Su Yadana,<sup>1</sup> Kristen Kelli Coleman,<sup>1</sup> Tham Thi Nguyen,<sup>1</sup> Christophe Hansen-Estruch,<sup>2</sup> Shirin Kalimuddin,<sup>3</sup> Koh Cheng Thoon,<sup>4</sup> Jenny Guek Hong Low,<sup>1,3</sup> Gregory Charles Gray<sup>1,5,6</sup>

<sup>1</sup>Program in Emerging Infectious Diseases, Duke-NUS Medical School, Singapore; <sup>2</sup>Duke University School of Medicine, Durham, USA; <sup>3</sup>Department of Infectious Diseases, Singapore General Hospital, Singapore; <sup>4</sup>Department of Pediatrics, Infectious Disease Service, KK Women's and Children's Hospital, Singapore; <sup>5</sup>Division of Infectious Diseases, School of Medicine and Global Health Institute, Duke University, Durham, NC, USA; <sup>6</sup>Global Health Research Center, Duke Kunshan University, Kunshan, China

## Significance for public health

We demonstrated the potential for airborne respiratory viruses to circulate among hospitalized children, nursing staff and visitors. We argue that bioaerosol sampling could serve as a noninvasive and low-cost method to monitor for novel respiratory virus incursions in clinical settings, and better understand the risk of acquiring a respiratory illness during a hospital visit.

sion through the inhalation of virus-laden particles. Specifically, airborne virus-laden particles  $\leq 4\mu\text{m}$  are thought to play a significant role in respiratory virus transmission as they can remain in the air for prolonged periods of time and are inhaled deep into the lungs.<sup>2</sup>

## Abstract

There is an increasing body of evidence suggesting that transmission of respiratory viruses occurs through the inhalation of virus-laden particles. Our study describes the use of an aerosol sampling system to monitor the prevalence of airborne viruses in a hospital setting. Using SKC AirCheck Touch pumps, with National Institute for Occupational Safety and Health (NIOSH) bioaerosol samplers and SKC filter cassette blanks, 28 aerosol samples were collected in a hospital ward in Singapore. Following DNA/RNA extraction, real-time RT-PCR/PCR was used for the detection of influenza A, B and D viruses, coronaviruses, enteroviruses, and adenoviruses. Airborne virus was detected in nine (32%) of 28 samples. Among the nine positive samples, eight were PCR-positive for adenovirus and one for influenza A virus. Our data suggest that bioaerosol sampling could be valuable in monitoring for airborne respiratory viruses in clinical environments to better understand the risk of infection during a hospital visit.

## Introduction

Despite efforts to reduce the incidence of nosocomial infections by measures such as practicing good hand hygiene and the use of personal protective equipment, hospital-acquired infections are still frequent. Two of the largest hospitals in Singapore report that on average, one in seven hospitalized patients acquire a nosocomial infection,<sup>1</sup> with immunocompromised children at greatest risk. Little is known regarding the transmission of respiratory viruses in clinical environments. Theoretically, influenza and other respiratory viruses are transmitted through contact with contagious persons and contaminated fomites. However, there is an increasing body of evidence supporting the concept of transmis-

## Materials and Methods

### Aerosol sampling

Aerosol samples were collected in a general pediatric ward at KK Women's and Children's Hospital, Singapore, using three National Institute for Occupational Safety and Health (NIOSH) two-stage cyclone samplers and one SKC filter cassette preloaded with a 37mm polytetrafluoroethylene (PTFE) filter (0.3 $\mu\text{m}$  pore size) designed to sample for severe acute respiratory syndrome-associated coronavirus (SARS-CoV). Aerosol samples were collected once per week for seven weeks in May and June 2017. The NIOSH samplers were stationed on tripods and placed along the corridor outside the open patient bedding area, and one SKC filter cassette was attached to a mobile computer on wheels (COW) used by doctors and nurses during ward rounds. Each sampler was connected to an AirChek® TOUCH Sample Pump (SKC, Eighty-Four, USA) with Tygon tubing (61 cm length, 0.635 cm diameter) for air collection at a rate of 3.5 L/min. A total of 840 L of air was collected during each four-hour sampling period. Each NIOSH sampler separates collected particles into three aerodynamic diameters:  $>4\mu\text{m}$ , 1-4 $\mu\text{m}$ , and  $<1\mu\text{m}$ .<sup>3</sup> Filter cassettes and sample tubes from the NIOSH samplers were stored at  $-80^\circ\text{C}$  before processing. Prior to nucleic acid extraction, sample material collected in the 1-4  $\mu\text{m}$  and  $<1\mu\text{m}$  size fractions were combined (described below).

### Nucleic acid extraction

PTFE filters were removed from the cassettes attached to the NIOSH samplers, transferred to 50 mL falcon tubes and vortexed for 15 s. One mL of 0.5% bovine serum albumin (BSA) fraction V in molecular grade water was then added to each 50mL falcon tube and vortexed again for 15 s. One mL of 0.5% BSA was added to each 1.5mL conical tube from the NIOSH samplers and vortexed for 15 s. These BSA solutions were then pooled into a 2mL cryovial tube. Two mL of 0.5% BSA fraction V solution was added to each 15mL falcon tube from the NIOSH samplers, vor-

texted for 15 s, and transferred to a cryovial tube and stored at -80°C until further used. Styrene filters from SKC cassettes were swabbed with FLOQSwabs soaked in 0.5% BSA fraction V solution. Swabs were then placed in 50mL falcon tubes, vortexed for 15 s, and transferred to cryovials. QIAamp viral RNA kit and QIAamp DNA Blood kit (Qiagen) were then used to extract RNA and DNA, respectively, from the sample solutions following the manufacturer's protocol.

### Real-time RT-PCR/PCR

RNA was tested for influenza A, B, and D,<sup>4-6</sup> coronavirus,<sup>7</sup> and enterovirus<sup>8</sup> using Superscript III One-step RT-PCR with Platinum Taq Polymerase. Extracted DNA was tested for adenovirus by real-time PCR using a QuantiNova Probe PCR kit (Qiagen) (Table 1).<sup>8</sup>

### Cell culture

For adenovirus-positive aerosol samples, 500 µL of sample was inoculated into adenocarcinomic human alveolar basal epithelial (A549) cells (ATCC® CCL185™) with Dulbecco's Modified Eagle Medium (DMEM) 2% (v/v) Fetal Bovine Serum (FBS), and incubated at 37°C. After 72 hours, inoculated shell vials were observed for cytopathic effect (CPE) daily for ten days. For influenza A virus-positive aerosol samples, 200 µL of sample was inoculated into Madin Darby Canine Kidney (MDCK) cells (ATCC® PTA-6500™) with DMEM containing 100 U/mL penicillin, 100 µg/mL streptomycin, 0.2% (w/v) BSA, 25 mM 4-(2-hydroxyethyl)-1-piperazineethanesulfonic acid (HEPES) buffer, and 1 µg/mL Tosyl phenylalanyl chloromethyl ketone (TPCK)-treated trypsin, and incubated at 37°C for 7 days with daily observations for CPE.

## Results

Eight (28.5%) of the 28 aerosol samples tested positive for adenovirus and one (3.5%) tested positive for influenza A virus. Aerosol samples with real-time RT-PCR/PCR cycle threshold (Ct) values <40 were considered positive. All eight adenovirus-positive samples were retrieved from the NIOSH samplers, 3 (37.5%) of which were from particles >4µm in aerodynamic diameter, and 5 (62.5%) from particles ≤4µm in aerodynamic diameter. The captured influenza A virus-positive particles were retrieved from a mobile SKC filter cassette and were therefore ≥0.3µm. None of the aerosol samples tested positive for influenza B or D virus, enterovirus or coronavirus. Attempts to grow viruses in cell culture from positive aerosol samples were unsuccessful.

## Discussion

Our pilot study provides molecular evidence of airborne respiratory viruses in a general pediatric ward in Singapore. Our results illustrate the potential of airborne respiratory viruses to circulate among hospitalized children, nursing staff and visitors. Additionally, we detected respirable virus-laden particles (≤4µm in diameter) which are thought to play a significant role in respiratory virus transmission. Aerosol samples testing positive for influenza A virus and adenovirus demonstrates the potential of these viruses to be transmitted via the airborne route. However, we were unable to document the viability of the virus particles and subsequent risk of infection. Additionally, the source of these viruses in the ward is

**Table 1. Sequences of primers and probes for RT-PCR/PCR assays.**

Assay	Primer Name	Primer Sequence (5'→3')
Influenza A Real-time	InfA-F Primer	GACCRATCCTGTACCTCTGAC
	InfA-R Primer	AGGGCATTYTGACAAAKCGTCTA
	InfA-P Probe	<b>FAM-TGCAGTCCTCGCTCACTGGGCACG-BHQ 1</b>
Influenza B Real-time	InfB-F Primer	TCCTCAAYTCACTCTTCGAGCG
	InfB-R Primer	CGGTGCTCTTGACCAAATTGG
	InfB-P Probe	<b>FAM-CCAATTCGAGCAGCTGAAACTGCGGTG-BHQ 1</b>
Influenza D Real-time	qRT-FluD-F	GCTGTTTGCAAGTTGATGGG
	qRT-FluD-R	TGAAAGCAGGTAACCTCAAGG
	qRT-FluD-Probe	<b>FAM-TTCAGGCAAGCACCCGTAGGATT-BHQ 1</b>
Coronavirus Real-time	Cov-NL63-F Primer	GTTCTGATAAGGCCACCATATAGG
	Cov-NL63-R Primer	TTTAGGAGGCAAAATCAACACG
	Cov-NL63-Probe	<b>TXR-CGCATACGCCAACGCTCTTGAACA</b>
	Cov-OC43-F Primer	CATACTCTGACGGTCACAATAATA
	Cov-OC43-R Primer	ACCTTAGCAACAGTCATATAAGC
	Cov-OC43-Probe	<b>YAK-TGCCCAAGAATAGCCAGTACCTAGT</b>
	Cov-HKU1-F Primer	TCCTACTAYTCAAGAAGCTATCC
	Cov-HKU1-R Primer	AATGAACGATTATTGGGTCCAC
	Cov-HKU1-Probe	<b>CY5-TYCGCCTGGTACGATTTTGCCTCA</b>
Adenovirus Real-time	Cov-229E-F Primer	CATACTATCAACCCATTCAACAAG
	Cov-229E-R Primer	CACGGCAACTGTCATGTATT
	Cov-229E-Probe	<b>FAM-ATGAACCTGAACACCTGAAGCCAATCTATG</b>
	ADVF	CAGGACGTCYTCGGAGTACCTGA
	ADVRI	CGGTGGTCACATCGTGGGT
Enterovirus Real-time	ADVRII	GCTGAAGTACGTVTCGGTGGC
	ADVRIII	GGTGAAGTAGGTGTCCGTGGC
	ADV Probe	<b>FAM-TGGTGCAGTTYGCCCG-BHQ 1</b>
	AN350-F Primer	GGCCCCTGAATGCGGCTAATCC
	AN351-R Primer	GCGATTGTACCATWAGCAGYCA
	AN234-Probe	<b>FAM-CCGACTACTTTGGGWTCCCGTGT-BHQ 1</b>

unknown as we did not recruit human subjects nor have access to patient records.

Despite previous successful attempts at using the NIOSH two-stage sampler and PCR analyses to collect and detect aerosolized influenza A virus,<sup>9,10</sup> all samples collected using the NIOSH sampler in this study tested negative for influenza A virus. We did capture and detect aerosolized adenovirus using the NIOSH samplers, however, none of the samples from the SKC filter cassettes tested positive for adenovirus. This result is inconsistent with our previous study which successfully recovered adenovirus DNA from aerosol samples collected using the SKC filter cassettes in patient waiting rooms.<sup>11</sup> Sampling sessions in our current study collected 840 L of air using one SKC filter cassette compared to the collection of 900 L of air per each of two SKC filter cassettes in our previous study. Additionally, SKC filter cassettes were mobilized in our current study and stationary in our previous study. The higher sample volumes collected in our previous study as well as the difference in mobility and location of the samplers might explain the difference in positive sample collections among studies.

In addition to bioaerosol sample collection and handling methods, environmental conditions can also influence the viability of airborne viruses and downstream virus recovery. For example, prolonged sampling periods can compromise stability of virus-laden aerosols and result in decreased viral recovery.<sup>12</sup> Additionally, it has been demonstrated that the survival of airborne influenza virus depends on ambient temperature, relative humidity (RH) and ultraviolet radiation levels.<sup>13</sup> Specifically, infectivity of influenza in a simulated examination room was reported to be the highest at 7–23% RH, moderate at 57% RH and lowest at 43% RH.<sup>14</sup> Although we did not record temperature and RH at our sampling site, a previous bioaerosol study recorded levels ranging from 54% to 68% RH in three different hospitals in Singapore.<sup>11</sup> High RH levels in Singapore could explain the low percentage of influenza A virus-positive aerosol samples in our study.

One strength of our bioaerosol sampling method is that it is a quick and non-invasive way to monitor for respiratory viruses without interrupting patients or healthcare professionals. Also, it requires little manpower to collect samples and results can be analyzed within a few hours. However, one limitation of our detection method was that we did not measure viral load in our aerosol samples, which makes it difficult to compare our results with quantitative aerosol studies in clinical settings. Additionally, our pilot study was not designed to collect patient data and therefore we were not able to match the virus-positive aerosol samples with individual patients present in the hospital ward at the time of sampling.

## Conclusions

In summary, we conducted a 7-week pilot study to monitor for aerosolized respiratory viruses among inpatients in a pediatric ward in Singapore. We found molecular evidence of influenza A virus and adenovirus, demonstrating the potential for airborne transmission. To comprehend this potential risk of transmission, our proof-of-concept project might be expanded in the future to specifically study patients with known positive clinical infections (e.g. through nasopharyngeal swabs) and determine how far away viable viruses can be detected from a patient's bedside. Additionally, future studies might involve more comprehensive demographic and clinical risk factor analyses to help us better understand phenomenon such as super-spreading. Lastly, bioaerosol surveillance might be useful in monitoring clinical populations for incursions of novel respiratory viruses, especially

since aerosol sampling in shared clinical spaces often requires no informed consent.

Correspondence: Kristen Kelli Coleman, Emerging Infectious Diseases, Duke-NUS Medical School, 8 College Road, 169857, Singapore.

Tel.: +65.66013667.

E-mail: kristen.coleman@duke-nus.edu.sg

Key words: respiratory viruses, aerosol, surveillance, adenovirus, influenza virus.

Acknowledgements: We thank Dr. Bill Lindsley from the National Institute for Occupational Safety and Health in the USA for loaning our team the bioaerosol samplers and guiding us in their use.

Contributions: SK, KT, JL, and GG designed the study. KT served as the sample site PI. Under KC's guidance, TN and CH collected the aerosol samples and TN, CH and SY performed laboratory works. TN, CH and SY analyzed the results. SY, KC and GG wrote the manuscript and TN helped format it.

Conflict of interest: The authors declare no conflict of interest.

Funding: This work was funded by SingHealth Foundation Research Grant [SHF/FG600P/2015].

Received for publication: 18 May 2018.

Revision received: 30 November 2018.

Accepted for publication: 20 December 2018.

©Copyright: the Author(s), 2019

Licensee PAGEPress, Italy

*Journal of Public Health Research* 2019;8:1407

doi:10.4081/jphr.2019.1407

This work is licensed under a Creative Commons Attribution NonCommercial 4.0 License (CC BY-NC 4.0).

## References

1. Singapore General Hospital. The New Paper (February, 2013). Retrieved from <https://www.sgh.com.sg/about-us/newsroom/News-Articles-Reports/Pages/hospital-acquired-infections.aspx>.
2. WHO. Infection prevention and control measures for acute respiratory infections in healthcare settings: An update. 2007:39–47.
3. Cao G, Noti JD, Blachere FM, et al. Development of an improved methodology to detect infectious airborne influenza virus using the NIOSH bioaerosol sampler. *J Environ Monitoring* 2011;13:3321–8.
4. World Health Organization (WHO). CDC protocol of realtime RT-PCR for influenza H1N1. World Health Organization, Geneva: Switzerland.
5. Selvaraju SB, Selvarangan R. Evaluation of three influenza A and B real-time reverse transcription-PCR assays and a new 2009 H1N1 assay for detection of influenza viruses. *J Clin Microbiol* 2010;48:3870–5.
6. Hause BM, Ducatez M, Collin EA, et al. Isolation of a novel swine influenza virus from Oklahoma in 2011 which is distantly related to human influenza C viruses. *PLoS Pathog* 2013;9:e1003176.
7. Vlasova AN, Halpin R, Wang S, et al. Molecular characterization of a new species in the genus Alphacoronavirus associated with mink epizootic catarrhal gastroenteritis. *J Gen Virol* 2011;92:1369–79.
8. Kilpatrick DR, Yang CF, Ching K, et al. Rapid group-, serotype-, and vaccine strain-specific identification of

- poliovirus isolates by real-time reverse transcription-PCR using degenerate primers and probes containing deoxyinosine residues. *J Clin Microbiol* 2009;47:1939-41.
9. Blachere FM, Lindsley WG, Pearce TA, et al. Measurement of airborne influenza virus in a hospital emergency department. *Clin Infect Dis* 2009;48:438-40.
  10. Coleman, K. Environmental Detection and Quantification of Airborne Influenza A Virus in an Elementary School, and its Implications for Student and Community Illness. (Electronic Thesis or Dissertation). 2017. Retrieved from <https://etd.ohiolink.edu/>.
  11. Nguyen TT, Poh MK, Low J, et al. Bioaerosol Sampling in Clinical Settings: A Promising, Noninvasive Approach for Detecting Respiratory Viruses. *Open Forum Infect Dis* 2017;4:ofw259.
  12. Blachere FM, Lindsley WG, Slaven JE, et al. Bioaerosol sampling for the detection of aerosolized influenza virus. *Influenza Other Respir Viruses* 2007;1:113-20.
  13. Cao G, Noti JD, Blachere FM, et al. Development of an improved methodology to detect infectious airborne influenza virus using the NIOSH bioaerosol sampler. *J Environ Monit* 2011;13:3321-8.
  14. Noti JD, Blachere FM, McMillen CM, et al. High humidity leads to loss of infectious influenza virus from simulated coughs. *PLoS One* 2013;8:e57485.

Non-commercial use only